


Entresto and Acute kidney failure - from FDA reports

On Jul, 23, 2017

9,385 people reported to have side effects when taking Entresto.

Among them, **140** people (1.49%) have Acute kidney failure

Number of reports submitted per year:



Time on Entresto when people have Acute kidney failure *:

- < 1 month: 53.66 %
- 1 - 6 months: 36.59 %
- 6 - 12 months: 9.76 %
- 1 - 2 years: 0.0 %
- 2 - 5 years: 0.0 %
- 5 - 10 years: 0.0 %
- 10+ years: 0.0 %

Gender of people who have Acute kidney failure when taking Entresto *:

- female: 22.12 %
- male: 77.88 %

Age of people who have Acute kidney failure when taking Entresto *:

- 0-1: 0.0 %
- 2-9: 0.0 %
- 10-19: 0.0 %
- 20-29: 0.0 %
- 30-39: 0.0 %
- 40-49: 8.2 %

- 50-59: 16.39 %
- 60+: 75.41 %

Top conditions involved for these people *:

- Cardiac Failure (37 people, 26.43%)
- Cardiac Failure Chronic (20 people, 14.29%)
- Left Ventricular Failure (11 people, 7.86%)
- Atrial Fibrillation/flutter (5 people, 3.57%)
- Cardiac Failure Congestive (4 people, 2.86%)

Top co-used drugs for these people *:

- Lasix (5 people, 3.57%)
- Coreg (5 people, 3.57%)
- Aldactone (5 people, 3.57%)
- Revatio (4 people, 2.86%)
- Eliquis (4 people, 2.86%)

Top other side effects for these people *:

- Hypotension (23 people, 16.43%)
- Cardiac Failure (21 people, 15.00%)
- Hyperkalemia (10 people, 7.14%)
- Death (10 people, 7.14%)
- Breathing Difficulty (9 people, 6.43%)

* Approximation only. Some reports may have incomplete information.

NOTE: The study is based on active ingredients and brand name. Other drugs that have the same active ingredients (e.g. generic drugs) are NOT considered.

WARNING: Please DO NOT STOP MEDICATIONS without first consulting a physician since doing so could be hazardous to your health.

DISCLAIMER: All material available on eHealthMe.com is for informational purposes only, and is not a substitute for medical advice, diagnosis, or treatment provided by a qualified healthcare provider. All information is observation-only, and has not been supported by scientific studies or clinical trials unless otherwise stated. Different individuals may respond to medication in different ways. Every effort has been made to ensure that all information is accurate, up-to-date, and complete, but no guarantee is made to that effect. The use of the eHealthMe site and its content is at your own risk.

You may report adverse side effects to the FDA at <http://www.fda.gov/medwatch/> or 1-800-FDA-1088 (1-800-332-1088).

If you use this eHealthMe study on publication, please acknowledge it with a citation: study title, URL, accessed date.
