



Cardiovascular manifestations and treatment considerations in covid-19

Yu Kang,¹ Tiffany Chen ^(D), ¹ David Mui,² Victor Ferrari,¹ Dinesh Jagasia,¹ Marielle Scherrer-Crosbie,¹ Yucheng Chen ^(D), ³ Yuchi Han ^(D)

ABSTRACT Since its recognition in December 2019, covid-19 has

INTRODUCTION

rapidly spread globally causing a pandemic. Pre-existing

severity and higher fatality rate of covid-19. Furthermore,

comorbidities such as hypertension, diabetes, and

cardiovascular disease are associated with a greater

covid-19 contributes to cardiovascular complications,

including acute myocardial injury as a result of acute

coronary syndrome, myocarditis, stress-cardiomyopathy,

arrhythmias, cardiogenic shock, and cardiac arrest. The

cardiovascular interactions of covid-19 have similarities

continued use of renin-angiotensin-aldosterone system

inhibitors, arrhythmia monitoring, immunosuppression or

Severe acute respiratory syndrome coronavirus 2

(SARS-CoV-2), which causes covid-19, was first

reported to WHO as a pneumonia of unknown

cause in Wuhan, China, on 31 December 2019.¹

While the initial outbreak was mostly confined

to the epicentre in China, SARS-CoV-2 quickly

spreads internationally causing a global pandemic. By 15 March, the number of cases outside of China

surpassed those in China, and as of 12 April 2020,

over 1.8 million cases and 110000 deaths were

many other viral infections.^{3 4} According to a study

by the Chinese Center for Disease Control and

Prevention of 44672 laboratory confirmed, 10567

clinically diagnosed and 16186 suspected cases of

covid-19, 81.4% exhibited mild illness (with no or

mild symptoms of pneumonia), 13.9% had severe

symptoms (dyspnoea with respiratory rate $\geq 30/$

min, SpO₂ ≤93%, PaO₂/FiO₂<300 and/or infiltra-

tion of lung field >50% within 24–48 hours), and

4.7% were critically ill (respiratory failure, septic

shock and/or multiorgan failure).⁵ In patients with

severe or critical disease, viral pneumonia can prog-

ress to acute respiratory distress syndrome (ARDS),

The most common symptoms of covid-19 include fever, dry cough, myalgia or fatigue, as is the case in

reported worldwide, affecting 185 countries.²

to that of severe acute respiratory syndrome, Middle

East respiratory syndrome and influenza. Specific

cardiovascular considerations are also necessary

in supportive treatment with anticoagulation, the

modulation, and mechanical circulatory support.

¹Cardiovascular Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania, USA ²University of Pennsylvania Perelman School of Medicine, Philadelphia, Pennsylvania, USA ³Department of Cardiology, West China Hospital, Sichuan University, Chengdu, China

Correspondence to

Dr Yuchi Han, Cardiovascular Medicine, University of Pennsylvania Perelman School of Medicine, Philadelphia, PA 19104, USA; yuchi.han@pennmedicine. upenn.edu

Received 4 April 2020 Revised 13 April 2020 Accepted 15 April 2020



© Author(s) (or their employer(s)) 2020. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Kang Y, Chen T, Mui D, et al. Heart Epub ahead of print: [please include Day Month Year]. doi:10.1136/ heartjnl-2020-317056

BMJ

and multisystem failure accompanied by a cytokine storm.⁵ Since patients with covid-19 with cardiovascular comorbidities have higher mortality,⁵ and the severity of covid-19 disease correlates with SARS-Co

Since patients with covid-19 with cardiovascular comorbidities have higher mortality,⁵ and the severity of covid-19 disease correlates with cardiovascular manifestations,⁶ it is important to understand the interaction of covid-19 and cardiovascular disease (CVD). This review will summarise our current understanding of the cardiovascular manifestations of covid-19, as compared with SARS (caused by SARS-CoV), the Middle East respiratory syndrome (MERS) (caused by MERS-CoV) and influenza. We will also discuss cardiovascular considerations regarding treatment strategies.

CARDIOVASCULAR COMORBIDITIES

The prevalence of diabetes mellitus (DM) and obesity in patients with covid-19 in the USA appear to be higher than in the general population, but the prevalence of CVD appears to be similar (table 1). However, in hospitalised patients with covid-19 with more severe disease, DM, hypertension and CVD seem to be more prevalent in both the USA and China, similar to MERS and influenza (table 2). A retrospective, single-centre case series of 187 patients with covid-19 found that patients with underlying CVD were more likely to have cardiac injury (troponin (Tn) elevation) compared with patients without CVD (54.5% vs 13.2%).6 In-hospital mortality was 7.6% for patients without underlying CVD and normal Tn, 13.3% for those with CVD and normal Tn, 37.5% for those without CVD but elevated Tn and 69.4% for those with CVD and elevated Tn.⁶ These observations suggest that although patients with pre-existing CVD may not be more susceptible to contracting SARS-CoV-2, they are prone to more severe complications of covid-19 with increased mortality.⁷ The association of cardiovascular and other comorbidities with case fatality rate is summarised in figure 1.

The higher mortality in patients with cardiovascular comorbidities may be directly attributable to underlying CVD or merely coincident with CVD. CVD may also be a marker of accelerated ageing, immune system dysregulation, or other mechanisms that affect covid-19 prognosis. Age is both a risk factor for CVD and an important determinant of covid-19 fatality, which has been found to be 14.8% in patients older than 80 years of age but <4% in patients younger than 70 years of age.⁵ Patients with elevated Tn levels were older than patients with normal Tn (71.4±9.4 vs 53.5±13.2 years).⁶ In addition, ageing weakens the immune system, as demonstrated by the lower protective titres after influenza vaccination in 50% of adults older than 65 years of age.⁸

PATHOPHYSIOLOGY OF CARDIAC INJURY

SARS-CoV-2 is an enveloped, positive-sense singlestranded RNA virus.⁹ SARS-CoV-2 and other similar coronaviruses use the ACE 2 (ACE2) protein for ligand binding before entering the cell via





Table 1	Prevalence of cardiopulmonary comorbidities in covid-19
disease ir	the USA and China compared with the general population

	Covid-19		General popula	ation
Geographic region	USA	China	USA (>20 years of age)	China (>15 years of age)
Total number	7162	44672	265 million	Various
Diabetes mellitus	10.9%	5.3%	9.8%	10.9%
Hypertension		12.8%	45.6%	23.2%
Cardiovascular disease	9.0%	4.2%	9.0%	3.7%*
Obesity	48.3%		31.2%	11.9%
Chronic lung disease	9.2%	2.4%	5.9%†	8.6%
Source (reference)	CDC ⁷⁴	Chinese CDC⁵	AHA ⁷⁵ Croft <i>et al</i> ⁷⁶	Hu et al ⁷⁷

*Combination of cerebrovascular disease, acute myocardial infarction, and heart failure.

†Chronic obstructive lung disease only.

AHA, American Heart Association; CDC, Center for Disease Control and Prevention.

receptor-mediated endocytosis.¹⁰ Recent data on viral structure reveal that SARS-CoV-2 has tighter interaction with the human ACE2 receptor binding domain as compared with SARS-CoV, which may explain in part the greater transmissibility of the current virus among humans.¹¹ ACE2 is a membrane protein that serves many physiological functions in the lungs, heart, kidneys and other organs.¹² It is highly expressed in type 2 lung alveolar cells, which provides an explanation for the respiratory symptoms experienced by patients with covid-19.¹³ More than 7.5% of myocardial cells have positive ACE2 expression, based on single-cell RNA sequencing,¹⁴ which could mediate SARS-CoV-2 entry into cardiomyocytes and cause direct cardiotoxicity.

The mechanisms of cardiovascular injury from covid-19 have not been fully elucidated and are likely multifactorial. SARS-CoV-2 viral particles have been identified by RT-PCR in cardiac tissue in some cases, ¹⁵ supporting that direct cardiotoxicity may occur. Virally driven hyperinflammation with cytokine release may lead to vascular inflammation, plaque instability, myocardial inflammation, a hypercoagulable state, and direct myocardial suppression.¹⁶ ¹⁷ Other systemic consequences of covid-19 infection, including sepsis and disseminated intravascular coagulation (DIC), may also mediate cardiac injury. Based on postmortem biopsies, the pathological features of covid-19 in multiple organs greatly resemble those seen in SARS^{18 19} and MERS.^{20 21} In cardiac tissue, pathological findings vary from minimal change to interstitial inflammatory infiltration and

myocyte necrosis (table 3). In the vasculature, microthrombosis and vascular inflammation could be found (table 3).

CARDIOVASCULAR MANIFESTATIONS

The clinical cardiovascular manifestations of covid-19 include elevation of cardiac biomarkers (ischaemic or non-ischaemic aetiology), cardiac arrhythmia, arterial and venous thromboembolism (VTE), and cardiogenic shock and arrest. The possible mechanisms and cardiovascular manifestations are shown in figure 2.

Elevation of cardiac biomarkers

Myocardial injury is common among patients with covid-19 infection and correlates with disease severity. Although somewhat variable, studies on patients with covid-19 have generally defined myocardial injury as the elevation of high-sensitivity cardiac troponin (hs-cTn) above the 99th percentile of its upper limit of normal or evidence of new electrocardiographic or echocardiographic abnormalities.^{3 22} Increased levels of hs-cTn correlate with disease severity and mortality rate in covid-19, even after controlling for other comorbidities.²³ A summary of studies with prevalence of myocardial injury and risk of severe disease or mortality is shown in table 4.

The pattern in the rise of cTn levels is also significant from a prognostic standpoint. Non-survivors had a higher level of Tn elevation which continued to rise until death (mean time from symptom onset to death was 18.5 days (IQR 15–20 days)), while Tn levels for survivors remained unchanged.²² This finding may support the monitoring of cTn levels every few days in hospitalised patients.

It is a challenging task to differentiate potential aetiologies of cardiac injury: acute coronary syndrome (ACS) due to plaque rupture or thrombosis (type I myocardial infarction (MI)) or supply-demand mismatch (type II MI), myocardial injury due to DIC, and non-ischaemic injury (myocarditis, stress-induced cardiomyopathy, or cytokine release syndrome).

Ischaemic myocardial injury *Myocardial infarction*

Severe viral infections can cause a systemic inflammatory response syndrome that increases the risk of plaque rupture and thrombus formation, resulting in either an ST-elevation MI or non-ST-elevation MI.²⁴ In a study of 75 patients hospitalised with SARS, acute MI was the cause of death in two of five fatal cases.²⁵ There is also a significant association between acute MI and influenza. As compared with the prevalence of acute MI

Table 2 Prevalence of comorbidities in hospitalised patients with severe respiratory viral infections									
	Covid-19			Influenza	SARS	MERS			
Geographic region	USA	China	Italy*	USA	Canada	Middle East, Europe			
Total number	178	416	1043	3271	144	637			
Diabetes mellitus	28.3%	14%	17.3%		11%	51%			
Hypertension	49.7%	30.5%	48.8%			49%			
Cardiovascular disease	27.8%	14.7%	21.4%	45.6%	8%	31%			
Obesity	48.3%			39.3%					
Chronic lung disease	34.6%	2.9%	4.0%	34.5%	1%				
Malignancy or immunocompromised	9.6%	2.2%	7.8%	17.0%	6%				
Source (reference)	CDC COVID-Net, Garg et al ⁷⁸	Shi <i>et al⁷</i>	Grasselli <i>et al</i> ⁷⁹	CDC FluServ-Net ⁸⁰	Booth <i>et al⁸¹</i>	Badawi and Ryoo ⁷³			

*Only including patients in the intensive care unit.

CDC, Centre for Disease Control and Prevention; MERS, Middle East respiratory syndrome; SARS, severe acute respiratory syndrome.

	Covid-19	Influenza	SARS	MERS
				- - -
Geographic region	China	Worldwide	Hong Kong	Middle East, Europe
Number of patients	44,672	610,782	115	637
Overall CFR	2.3%	0.1%	10%	35%
Without risk factors				
Diabetes mellitus	OR 9.07	OR 0.59	HR 4.6	
Hypertension	OR 7.39	OR 3.53		
Cardiovascular disease	OR 13.64	OR 1.97	HR 9.2	
Obesity		OR 2.74		
Chronic lung disease	OR 7.74	OR 1.80		
Cancer	OR 6.88	OR 2.67		
Source	Chinese CDC	Mertz <i>et al.</i>	Chan <i>et al.</i>	Badawi <i>et al.</i>

Influence of Comorbidities on Case Fatality Rate (CFR)

Figure 1 Influence of cardiovascular and other comorbidities on CFR from respiratory viral infections. CDC, Center for Disease Control and Prevention; CFR, case fatality rate; MERS, Middle East respiratory syndrome; SARS, severe acute respiratory syndrome. Data are from Chinese CDC,⁵ Mertz *et al*,⁷¹ Chan *et al*,⁷² and Badawi *et al*.⁷³

occurred 1 year before or 7 days to 1 year after influenza, the incidence ratio of acute MI within 7 days of influenza infection was 6.1 (95% CI: 3.9 to 9.5).²⁶ Although reports of type I MI in patients with covid-19 have not yet been published, anecdotal report was presented.²⁷ Treatment of ACS in covid-19 should be according to the updated Society for Cardiovascular Angiography and Interventions guidelines.²⁸

Severe respiratory viral infections can also lead to decreased oxygen delivery to the myocardium via hypoxaemia and vasoconstriction, as well as the haemodynamic effects of sepsis with increased myocardial oxygen demand. This supply and demand mismatch may lead to sustained myocardial ischaemia in patients with underlying coronary artery disease. However, a rise and/ or fall of hs-cTn is not sufficient to secure the diagnosis of acute MI as seen in MI with non-obstructive coronaries, even in the absence of covid-19. Therefore, the diagnosis of acute MI should also be based on clinical judgement, symptom/signs, ECG changes, and imaging studies.

Myocardial injury with DIC

DIC is a life-threatening condition present in 71.4% (15/21) of non-survivors with covid-19 and 0.6% (1/162) of survivors.²⁹ A marker of severe sepsis, DIC further perpetuates multiorgan damage through thrombosis, reduced perfusion, and bleeding, DIC has been implicated in the thrombosis of coronary arteries (epicardial vessels and microvasculature), focal necrosis of the myocardium, and severe cardiac dysfunction.³⁰ Myocardial injury with DIC has been recently reported in two critically ill patients with covid-19.³¹ Both patients had significantly elevated Tn and brain natriuretic peptide, which normalised after treatment with heparin, mechanical ventilation, and antiviral agents.

Non-ischaemic myocardial injury Myocarditis and stress-induced cardiomyopathy

Myocardial injury from SARS-CoV-2 infection may also be mediated by non-ischaemic mechanisms, such as acute and fulminant myocarditis and stress-induced cardiomyopathy. Reports of various presentations of non-ischaemic myocardial injury in covid-19 are summarised in table 5. The distinction between myocarditis and stress-induced cardiomyopathy can be challenging, since cardiovascular magnetic resonance (CMR) and/or biopsy are not available in most cases. Fried et al and Sala et al each reported a patient with covid-19 with mid-left ventricular (LV) or basal-to-mid LV hypokinesis, a pattern of mid-ventricular, or reverse Takotsubo stress cardiomyopathy, respectively.^{32 33} The incidence of acute heart failure was 33% (7/21) in critically ill patients with covid-19 without a past history of LV systolic dysfunction in Washington state.³⁴ Importantly, cardiomyopathy can develop in covid-19 with mild or absent respiratory symptoms.35

Myocardial injury with cytokine release syndrome

Similar to SARS-CoV and MERS-CoV, SARS-CoV-2 can elicit the intense release of multiple cytokines and chemokines by the immune system.^{3 36} Cytokine release syndrome (aka 'cytokine storm'), a poorly understood immunopathological process caused by hyperinduction of proinflammatory cytokines such as interleukin (IL)-1, IL-6, T helper 1 cytokine interferon-gamma, and tumour necrosis factor-alpha (TNF- α), has been reported in the setting of SARS, MERS, and influenza.^{37–39} It is postulated that proinflammatory cytokines depress myocardial function immediately through activation of the neural sphingomyelinase

Table 3 F	3 Pathological findings of the cardiopulmonary systems in death related to coronaviruses						
Study	Region	Disease	Age/Sex	Lung	Heart	Vasculature	
Yao <i>et al</i> ¹⁵	China	Covid-19	63/M, 69/M, 79/F	Alveolar exudates, interstitial inflammatory infiltration and fibrosis, hyaline membrane formation. Positive RT-PCR for SARS-CoV-2, positive viral inclusions.	Myocardial hypertrophy and multifocal necrosis, interstitial inflammatory infiltration. No viral inclusions; negative RT-PCR for SARS-CoV-2.	Diffused hyaline thrombosis in microcirculation in multiple organs.	
Xu et al ⁸²	China	Covid-19	50/M	Bilateral diffuse alveolar damage with cellular fibromyxoid exudates, desquamation of pneumocytes and hyaline membrane formation, pulmonary oedema, interstitial mononuclear inflammatory infiltration.	A few interstitial inflammatory infiltrations.	NA	
Tian <i>et al⁸³</i>	China	Covid-19	78/F, 74/M, 81/M, 59/M	Diffused alveolar damage, hyaline membrane formation and vascular congestion, inflammatory cellular infiltration, focal interstitial thickening (case 3), large area of intra-alveolar haemorrhage and intra-alveolar fibrin cluster formation (case 4). Positive RT-PCR assay for SARS-CoV-2 in 1/3 patients.	Various degrees of focal oedema, interstitial fibrosis and myocardial hypertrophy; no inflammatory infiltration. Positive RT-PCR assay for SARS-CoV-2 in 1/2 patients (both patients with elevated troponin).	Fibrinoid necrosis of the small vessels of lung (case 4).	
Fox <i>et al⁸⁴</i>	USA	Covid-19	Four patients, range: 44—76 years	Pleural effusion, bilateral pulmonary oedema, patches of haemorrhage, diffuse alveolar damage, lymphocytic infiltration, hyaline membrane and fibrin. Viral inclusion.	Pericardial effusion, cardiomegaly with RV dilatation, scattered individual myocyte necrosis.	Thrombi and lymphocytic infiltration of small vessels of lung.	
Lang <i>et al^{es}</i>	China	SARS	73/M, 64/F, 69/F	Oedema and homogeneous fibrinous deposition of alveolar walls, desquamation of pneumocyte; exudate in alveolar space, hyaline membrane formation; inflammatory infiltration. Positive RT-PCR assay for SARS-CoV-1 in all three patients.	Atrophy of cardiac muscle, lipofuscin deposition in cytoplasm, proliferation of interstitial cells and lymphocytes.	Fibrous thrombi in pulmonary vessels.	
Ding <i>et al</i> ¹⁸	China	SARS	62/F, 25/M, 57/M	Extensive consolidation, pulmonary oedema, haemorrhagic infarction, desquamative alveolitis and bronchitis, exudation, hyaline membrane formation, focal necrosis. Viral inclusion.	Myocardial stromal oedema, inflammatory infiltration, hyaline degeneration and lysis of cardiac muscle.	Diffused inflammatory infiltration of vessel walls, edematous endothelial cells, fibrinoid necrosis of veins in multiple organs, mixed thrombi in small veins and hyaline thrombi in microvessels.	
Farcas <i>et al</i> ¹⁹	Canada	SARS	21 patients	Diffuse alveolar damage. Positive RT-PCR assay for SARS-CoV-1 in 9/13 patients.	Positive RT-PCR assay for SARS-CoV-1 in 7/18 patients.	NA	
Ng <i>et al</i> ²¹	UAE	MERS	45/M	Diffuse alveolar damage with denuding of bronchiolar epithelium, prominent hyaline membranes, alveolar fibrin deposits.	Diffuse myocyte hypertrophy, patchy fibrosis.	NA	
Alsaad <i>et al</i> ²⁰	Saudi Arabia	MERS	33/M	Hyaline membrane formation, diffuse alveolar damage, parenchymal necrosis. Viral inclusion.	No significant inflammatory infiltration.	Subendothelial inflammatory infiltration of interstitial arteries of the lung.	

MERS, Middle East respiratory syndrome; NA, not available; SARS, severe acute respiratory syndrome.

pathway and subacutely (hours to days) via nitric oxide-mediated blunting of beta-adrenergic signalling.⁴⁰ Accumulating evidence suggest that a subgroup of patients with severe covid-19 can develop cytokine storm.³⁶ Plasma levels of IL-1 β , IL-6, IL-8 and TNF- α have been found to be significantly higher in patients with covid-19.³ The clinical and biochemical profiles of non-survivors in patients with covid-19 with highly elevated ferritin and IL-6 also suggest that cytokine release contribute to mortality.⁴¹

Arrhythmia

Arrhythmia could be the first presentation of covid-19, and new-onset and/or progressive arrhythmia could indicate cardiac involvement. A study of 137 patients in Wuhan showed that 7.3% had experienced palpitations as one of their presenting symptoms for covid-19.⁴² The prevalence of unspecified arrhythmia in two additional studies is listed in table 4. Arrhythmias were

found to be more common in intensive care unit (ICU) patients with covid-19 (44.4%) than non-ICU patients (6.9%).⁴ Patients with elevated Tn also had a higher incidence of malignant arrhythmia (haemodynamically unstable ventricular tachycardia or ventricular fibrillation) than those with normal Tn levels $(11.5\% \text{ vs } 5.2\%, \text{ p} < 0.001).^6$

Venous thromboembolism

Due to prolonged immobilisation, hypercoagulable status, active inflammation, and propensity for DIC, patients with covid-19 are at increased risk of VTE. The prevalence of ultrasound-confirmed deep venous thrombosis in patients with covid-19 is 22.7%⁴³ and 27% in ICU patients.⁴⁴ Patients with covid-19 have been shown to have significant higher level of D-dimer, fibrin degradation products (FDP), and fibrinogen, compared with healthy controls.⁴⁵ In addition, D-dimer and FDP titres

Possible Mechanisms of Cardiovascular Injury Due to Covid-19



Figure 2 Possible mechanisms of cardiovascular injury due to covid-19. DIC, disseminated intravascular coagulation; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2.

were higher in patients with severe covid-19 than those with milder disease.⁴⁵ A retrospective, multicentre cohort study demonstrated that D-dimer $>1 \mu g/mL$ on admission was associated with in-hospital death (OR: 18.4, 95% CI: 2.6 to 128.6, p=0.003).²² Thus, in the setting of critically ill covid-19 patients with clinical deterioration, VTE (including pulmonary embolism) should be considered.

TREATMENT CONSIDERATIONS

There is currently no proven therapy for covid-19, although many are under investigation. We focus our discussion here on supportive therapies and considerations of potential therapies relevant to the cardiovascular system, summarized in table 6.

Study	Region	Dates of study	Number	Age*	Male (%)	Tn elevation (%)	Arrhythmia (%)	Death (%)	OR of death or severe disease‡ with Tn elevation
Huang <i>et al³</i>	China	16 December 2019–2 January 2020	41 (with 17.1% censored)	49 [41–58]	73.2	12.2		14.6	12.0 [1.2 - 121.8] [‡]
Zhou <i>et al</i> ²²	China	29 December 2020–31 January 2020	191	56 [46–67]	62.3	16.6		28.3	80.1 [10.3 - 620.4]
Wang <i>et al</i> ⁴	China	1 January 2020–28 January 2020	138 (with 61.6% censored)	56 [42–68]	54.3	7.2§	16.7	4.3	14.3 [2.9 - 71.1] [‡]
Shi <i>et al⁷</i>	China	20 January 2020–10 February 2020	416 (with 76.7% censored)	64 (21–95)†	49.3	19.7		14.1	53.2 [11.4 - 248.0]
Guo <i>et al⁶</i>	China	23 January 2020–3 February 2020	187	59±15	48.7	27.8	16.7	23.0	15.1 [6.7 - 34.1]
Chen <i>et al⁸⁶</i>	China	January 2020–February 2020	150	59±16	56.0	14.7		7.3	28.3 [9.2 - 87.2]‡

†Expressed as median (range).

‡OR of severe disease with troponin elevation.

§ Acute myocardial injury defined as biomarker above the 99th percentile upper reference limit or new abnormalities in electrocardiography or echocardiography IQR, interquartile range; OR, odds ratio; SD, standard deviation; Tn, troponin.

	e	ed, LVEF	id, LVEF day 6	ed, LVEF day 10	ulated from ifter 7 days, emain ianical on	intubated	ged	Continued
	Outcom	Recovere 68%	Improve 44% on	2 Recovere 50% on	Decannu ECMO a' stable, re on mech ventilati	Remain on day 9	Discharg	
	Treatment	Antiviral, CRRT, corticosteroid, immunoglobulin, high- flow oxygen	Antiviral, corticosteroid, CQ, dobutamine, medical treatment for HF	IABP, dobutamine, HCC	 HCQ, invasive ventilation and VV ECMO, after LV function deterioration, change to VAV ECMO 	Broad-spectrum antibiotics, nitroglycerin, furosemide, mechanical ventilation vasopressor	MMF discontinued, HCQ, azithromycin, ceftriaxone	
	Dx	Myocarditis?	, Myopericarditis	Myopericarditis, with cardiogenic shock	Stress cardiomyopath	Decompensated heart failure	Myocarditis?	
	Biomarkers	Tnl, IL-6, BNP elevated ,	hs-TnT, NT-proBNF elevated	Thl elevated	TnT, IL-6, ferritin, CRP elevated	TnT, NT-proBNP, ferritin elevated	hs-TnT, IL-6, NT- proBNP, ferritin elevated	
	Other diagnostic studies	TTE: diffuse dyskinesia, LVEF: 32%, PAP: 44 mm Hg RV normal	CXR: normal; CMR: LVH, BIV hypokinesis LVEF: 35%, BiV myocardial oedema, diffuse LGE	CXR: normal Cath: non-obstructiv RHC: RA: 10 mm Hg, PCWP 21 mm Hg, CI: 1 L/min/m ² . TTE: LVH, LVEF 30%, severe hypokinetic RV.	CXR: bilateral pulmonary opacities TTE normal (before VV ECMOI), TTE: LVEF 20%–25%, akinesis of mid-LV segments; mildly reduced RV function	CXR: bibasilar opacities, vascular congestion TTE: severely reducec LV function	TTE: normal cardiac allograft function	
	ECG	Sinus tachycardia	Low voltage, diffuse ST- elevation, ST depression in V1 and aVR	Sinus tachycardia, low QRS voltage, diffuse ST and PR elevations, ST depression in aVR	SVT, Sinus tachycardia, AIVR	Sinus, PVC, PAC, lateral T inversion, QTc 528 ms	New T-wave inversion	
covid-19	Symptoms/Signs	Fever, cough, dyspnoea	Fever, dry cough, fatigue, hypotensive, normal oxygen saturation	Chest pressure, afebrile, no respiratory symptoms, normal oxygen saturation	Cough, chest pain, dyspnoea, rapidly deteriorated respiratory status	Non-productive cough, afebrile, dyspnoea, oxygen saturation 88%	Fever, dry cough, dyspnoea	
myocardial injury in	Comorbidities	Allergic cough	None	HTN, hyperlipidaemia	M	NICM with normal LVEF, AF, HTN, DM	Heart and renal transplant	
n-ischaemic	Sex	Σ	ш	ι.	Σ	њ.	Σ	
racteristics of no	Age	63	23	5	ŝ	64	51	
Clinical cha	Region	China	Italy	USA	USA	USA	USA	
Table 5 (Study	Zeng <i>et al⁸⁷</i>	Inciardi <i>et al^{3:}</i>	Fried <i>et al³²</i>	Fried <i>et al</i> ³²	Fried <i>et al³²</i>	Fried <i>et al³²</i>	

Heart: first published as 10.1136/heartjnl-2020-317056 on 30 April 2020. Downloaded from http://heart.bmj.com/ on May 2, 2020 by guest. Protected by copyright.

Table 5	Continued										
Study	Region	Age	Sex	Comorbidities	Symptoms/Signs	ECG	Other diagnostic studies	Biomarkers	X	Treatment	Outcome
Sala et af ^a AF, atrial fibh	Italy rillation, AIVR, acc	43 defrated idioventric	F cular rhythm	None 5 Biv, biventricular, BNP, E	Chest pain, dyspnoea, oxygen saturation 89% chance of the saturation 89% chan hatriuretic peptide; C	Sinus, new non- specific T-wave changes , cardiac index; CMI	CXR: multifocal bilateral opacities; Coronary CTA normal; 3D CT: mid-basal LV hypokinesia, normal apical function TTE: LVEF 43%, inferior wall hypokinesis; CMR on D7: LVEF 64%, mild hypokinesis at mid hypokinesia at mid myocardial oedema, no LGE; EMB: lymphocytic infiltration, interstitial oedema, interstitial oedema, inte	hs-TnT, NT-proBNP I elevated retronance; CPAP, e	Ayocarditis Ontinuous positive ai	CPAP, antiviral, HCQ way pressure; CQ, chloro	Discharged auine: CRP, C
reactive pro hydrochloro	tein; CRRT, continu quine; HF, heart fa	June: HTN, hyperter Lous renal replacen Jure; HTN, hyperter	nent therapy nsion; IABP, i al nrohrain n	; CTA, computed tomogra intra-aortic balloon pump patriuratic nenticle: PAC n	phy angiogram; CXR, ches phy angiogram; CXR, ches remature atrial complex. F	st X-ray; DM, diabete ancement; LVEF, left AP nulmonary arter	se mellitus; Dx, diagnos se mellitus; Dx, diagnos : ventricular ejection fra v nressure: PCWP nulm	s; ECMO, extracorport ction; LVH, left ventri onary canillary wedge	al membrane oxygen ular hypertrophy; MM nressure PVC nrema	ation; EMB, endomyocard IF, mycophenolate mofetil	ial biopsy; HCQ, ; NICM, non- RHC richt heart

restriction canonitypeous, we product we emined problem natureter peptud; rest, premate and compres, rest, pumorial areny pressure, rest, pumorial canonity areny pressure, rest, pumorial canonity of the second structure rest, rest, premated arend structure rest, premated arend pressure, rest, pumorial arend pressure, rest, pumorial canonity arend pressure, rest, pumorial arend pressure, rest, pumori

Heart: first published as 10.1136/heartjnl-2020-317056 on 30 April 2020. Downloaded from http://heart.bmj.com/ on May 2, 2020 by guest. Protected by copyright.

Table 6 Cardiovascular considerations in treatment	
Cardiovascular concerns	Treatment considerations
STEMI and NSTEMI	Primary PCI vs thrombolytics
Myocardial injury	Worse prognosis, monitoring rising trends
Hypercoaulable state	Thromboprophylaxis
ACEI or ARB use	Continue treatment currently, await further studies
HCQ, CQ and/or azithromycin use	QTc monitoring, avoid other QTc prolonging drugs
Immunosupression/Immunomodulation	Maybe helpful in selected patients with cytokine storm
MCS	IABP and VA ECMO might be used for support in cardiogenic shock

ACEI, ACE inhibitor; ARB, angiotensin receptor blocker; CQ, chloroquine; HCQ, hydroxychloroquine; IABP, intra-aortic balloon pump; MCS, mechanical circulatory support; NSTEMI, non-ST-elevation myocardial infarction; VA ECMO, venoarterial extracorporeal membrane oxygenation.

Anticoagulant therapy

Due to the high rate of associated arterial thromboembolism and VTE, prophylactic anticoagulation is essential in the management of hospitalised patients with covid-19,44 46 although the optimal thromboprophylaxis regimen is unclear. In a retrospective study of 449 patients with severe covid-19, 99 patients received unfractionated heparin or low molecular weight heparin for at least 7 days.⁴⁷ No difference in overall 28-day mortality was observed, but in subgroups of patients with sepsis-induced coagulopathy score ≥ 4 , or D-dimer > sixfold of upper limit of normal, the heparin group had lower mortality compared with the no-heparin group (40.0% vs 64.2%, p=0.029), (32.8% vs 52.4%, p=0.017), respectively.47 Another concern regarding thromboprophylaxis is the drug-drug interaction between some antiviral treatments (such as ribavirin, lopinavir and ritonavir) and direct oral anticoagulants. Low molecular weight heparin is likely preferred in critically ill patients with covid-19, if anticoagulation is used.

ACE inhibitors and angiotensin receptor blockers

Since SARS-CoV-2 uses the ACE2 protein for ligand binding before entering the cell,¹⁰ there are concerns regarding the use of renin-angiotensin-aldosterone system (RAAS) inhibitors that may increase ACE2 expression.^{13 48} The relationship between ACE2 expression and SARS-CoV-2 virulence is uncertain. After cell entry via ACE2, SARS-CoV-2 appears to subsequently downregulate ACE2 expression, which is vital to maintenance of cardiac function.⁴⁹ In mouse models, reduction in ACE2 expression is associated with increased severity of lung injury induced by influenza.⁵⁰ Whether higher expression of ACE2 increases susceptibility to SARS-CoV-2 infection or confers cardioprotection remains controversial.⁵¹ Furthermore, experimental animal models and few studies in humans have yielded conflicting results as to whether RAAS inhibition increases ACE2 expression.^{52 53} Based on the uncertainty regarding the overall effect of RAAS inhibitors (ACE inhibitor (ACEI) and angiotensin receptor blocker (ARB) therapy) in covid-19, multiple specialty societies currently recommend that RAAS inhibitors be continued in patients in otherwise stable condition.⁵¹

Hydroxychloroquine/Chloroquine and azithromycin

Chloroquine is a versatile bioactive agent with antiviral activity in vitro against DNA viruses as well as various RNA viruses, including SARS-CoV, MERS-CoV, and SARS-CoV-2, prompting it to be studied for patients with covid-19.⁵⁴ Similar to chloroquine, hydroxychloroquine confers antiviral effects and has an additional modulating effect on activated immune cells to decrease IL-6 expression.⁵⁵ Non-randomised, observational studies of >100 patients at 10 hospitals in China

suggest that chloroquine or hydroxychloroquine is superior to the control treatment in resolving pneumonia (based on clinical and imaging findings), promoting conversion to viral negative status and shortening the disease course.⁵⁶ In a small open-label, non-randomised study conducted in France, 70% of hydroxychloroquine-treated patients with covid-19 had undetectable viral titres at 6 days, compared with 12.5% in the control group (p=0.001),⁵⁷ although there were major methodological concerns with the study. Azithromycin, a macrolide antibiotic which acts against Zika and Ebola viruses in vitro⁵⁸ and suppresses inflammatory processes,⁵⁹ has been proposed as an effective adjunct to hydroxychloroquine in covid-19 through unclear mechanisms.⁵⁷ Although the preliminary evidence for quinoline antimalarials may be promising, it should be taken with caution until randomised clinical trials are completed.

Both chloroquine and azithromycin have generally favourable safety profiles, but they are known to cause cardiovascular side effects including prolongation of QT interval.⁶⁰ Although azithromycin interferes minimally with the cytochrome P-450 system in vitro,⁶¹ chloroquine is metabolised by CYP2C8 and CYP3A4/5,⁶² and there is a potentially enhanced risk of significant QT prolongation induced by combination therapy. Guidance for managing QTc prolongation in covid-19 pharmacotherapies and other cardiac electrophysiology issues related to covid-19 has recently been published by the Heart Rhythm Society.⁶³

Immunosuppressive therapy

As for SARS and MERS, corticosteroids are not routinely recommended for covid-19 and might exacerbate the resulting lung injury.⁶⁴ However, given the detrimental effects of the cytokine release syndrome associated with covid-19 on the cardiopulmonary system, immunosuppressive therapy to dampen the hyperinflammatory response may be beneficial.³⁶ Screening patients with covid-19 using laboratory data (increasing ferritin and erythrocyte sedimentation rate, and decreasing platelets) and a proposed score for the hemophagocytic response (HScore)⁶⁵ for hyperinflammation may help to identify patients for whom immunosuppression might improve survival.³⁶

IL-6 inhibitors may have a role as immunomodulators. Tocilizumab, an inhibitor of the IL-6 receptor, was shown to effectively improve symptoms and prevent clinical deterioration in a retrospective case series study of 21 covid-19 patients with severe or critical disease.⁶⁶ Several clinical trials are ongoing to assess the efficacy and safety of tocilizumab (NCT04317092)⁶⁷ or sarilumab, another IL-6 inhibitor (NCT04327388)⁶⁸ in patients with covid-19.

Mechanical cardiopulmonary support

Mechanical cardiopulmonary support in respiratory failure has been reported with variable survival rates. Extracorporeal Life Support Organization Registry is being adapted to investigate covid-19 in prospective observational studies addressing the role of extracorporeal membrane oxygenation (ECMO) in these critically ill patients.⁶⁹

In the setting of cardiogenic shock related to covid-19, intraaortic balloon pump (IABP), or veno-arterial ECMO should be considered. During the 2009 H1N1 pandemic in Japan, 9 out of 13 patients with fulminant myocarditis receiving emergent mechanical circulatory support (IABP and/or percutaneous cardiopulmonary support) survived, while all four patients with fulminant myocarditis treated without mechanical circulatory support died.⁷⁰ A case has been reported of a patient with covid-19 without respiratory symptoms presenting with cardiogenic shock attributed to acute myopericarditis and successfully treated with IABP support.³² Perhaps a more likely scenario is the conversion of veno-venous ECMO for ARDS to venoarterial-venous ECMO on development of cardiogenic shock, as described in another case report.³² Regardless of the mode of mechanical support, patient selection should involve consideration of comorbidities and potential complications.

CONCLUSION

Covid-19 is similar to SARS and MERS with regard to host vulnerability, specifically in those with substantial cardiovascular comorbidities. Greater transmissibility of covid-19 has resulted in a worldwide pandemic, a record number of infected individuals and an excess mortality that far exceeded previous coronavirus-related outbreaks. Myocardial injury is common in covid-19 and portends a worse prognosis. Differentiating between the various causes of myocardial injury is crucial to determining the treatment course. The cardiovascular considerations for treatment, including anticoagulation, ACEI or ARB use, anti-arrhythmic management, immunosuppression/modulation, and haemodynamic support, are important and continue to evolve.

Twitter Tiffany Chen @tiffchenMD

Contributors YK, TC, DM, and YH drafted the manuscript. All authors critically reviewed and edited the manuscript.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: http://creativecommons.org/licenses/by-nc/4.0/.

ORCID iDs

Tiffany Chen http://orcid.org/0000-0003-3768-3143 Yucheng Chen http://orcid.org/0000-0002-0601-8039 Yuchi Han http://orcid.org/0000-0001-7582-1848

REFERENCES

1 WHO. Novel coronavirus (2019-nCoV) situation Report-1, 2020. Available: https:// www.who.int/docs/default-source/coronaviruse/situation-reports/20200121-sitrep-1-2019-ncov.pdf?sfvrsn=20a99c10_4

- 2 JHU. Coronavirus COVID-19 global cases by the center for systems science AD engineering (CSSE) at Johns Hopkins University. Available: https://coronavirus.jhu.edu/ map.html [Accessed 12 Apr].
- 3 Huang C, Wang Y, Li X, *et al*. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 2020;395:497–506.
- 4 Wang D, Hu B, Hu C, et al. Clinical characteristics of 138 hospitalized patients with 2019 novel coronavirus-infected pneumonia in Wuhan, China. JAMA 2020;323:1061–9.
- 5 ChinaCDC. The epidemiological characteristics of an outbreak of 2019 novel coronavirus diseases (COVID-19) — China, 2020. *China CDC Weekly* 2020;2:113–22.
- 6 Guo T, Fan Y, Chen M, et al. Cardiovascular implications of fatal outcomes of patients with coronavirus disease 2019 (COVID-19). JAMA Cardiol 2020.
- 7 Shi S, Qin M, Shen B, et al. Association of cardiac injury with mortality in hospitalized patients with COVID-19 in Wuhan, China. JAMA Cardiol 2020. doi:10.1001/ jamacardio.2020.0950. [Epub ahead of print: 25 Mar 2020].
- 8 Govaert TM, Thijs CT, Masurel N, et al. The efficacy of influenza vaccination in elderly individuals. A randomized double-blind placebo-controlled trial. JAMA 1994;272:1661–5.
- 9 Zhou P, Yang X-L, Wang X-G, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature 2020;579:270–3.
- 10 Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor. Cell 2020. doi:10.1016/j.cell.2020.02.052. [Epub ahead of print: 04 Mar 2020].
- 11 Shang J, Ye G, Shi K, *et al.* Structural basis of receptor recognition by SARS-CoV-2. *Nature* 2020.
- 12 Zhang H, Penninger JM, Li Y, *et al*. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med* 2020;46:586–90.
- 13 Esler M, Esler D. Can angiotensin receptor-blocking drugs perhaps be harmful in the COVID-19 pandemic? J Hypertens 2020;38:781–2.
- 14 Zou X, Chen K, Zou J, et al. Single-cell RNA-seq data analysis on the receptor ACE2 expression reveals the potential risk of different human organs vulnerable to 2019nCoV infection. Front Med 2020. doi:10.1007/s11684-020-0754-0. [Epub ahead of print: 12 Mar 2020].
- 15 Yao XH, Li TY, He ZC, et al. [A pathological report of three COVID-19 cases by minimally invasive autopsies]. Zhonghua Bing Li Xue Za Zhi 2020;49:E009.
- 16 Prabhu SD. Cytokine-induced modulation of cardiac function. *Circ Res* 2004;95:1140–53.
- 17 Levi M, van der Poll T, Büller HR. Bidirectional relation between inflammation and coagulation. *Circulation* 2004;109:2698–704.
- 18 Ding Y, Wang H, Shen H, et al. The clinical pathology of severe acute respiratory syndrome (SARS): a report from China. J Pathol 2003;200:282–9.
- 19 Farcas GA, Poutanen SM, Mazzulli T, et al. Fatal severe acute respiratory syndrome is associated with multiorgan involvement by coronavirus. J Infect Dis 2005;191:193–7.
- 20 Alsaad KO, Hajeer AH, Al Balwi M, et al. Histopathology of Middle East respiratory syndrome coronovirus (MERS-CoV) infection - clinicopathological and ultrastructural study. *Histopathology* 2018;72:516–24.
- 21 Ng DL, Al Hosani F, Keating MK, et al. Clinicopathologic, immunohistochemical, and ultrastructural findings of a fatal case of middle East respiratory syndrome coronavirus infection in the United Arab Emirates, April 2014. Am J Pathol 2016;186:652–8.
- 22 Zhou F, Yu T, Du R, *et al*. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet* 2020;395:1054–62.
- 23 Zheng YY, YT M, Zhang JY, et al. COVID-19 and the cardiovascular system. Nat Rev Cardiol 2020.
- 24 Warren-Gash C, Hayward AC, Hemingway H, et al. Influenza infection and risk of acute myocardial infarction in England and Wales: a caliber self-controlled case series study. J Infect Dis 2012;206:1652–9.
- 25 Peiris JSM, Chu CM, Cheng VCC, et al. Clinical progression and viral load in a community outbreak of coronavirus-associated SARS pneumonia: a prospective study. Lancet 2003;361:1767–72.
- 26 Kwong JC, Schwartz KL, Campitelli MA, et al. Acute myocardial infarction after Laboratory-Confirmed influenza infection. N Engl J Med 2018;378:345–53.
- 27 ACC. ACC/Chinese cardiovascular association COVID-19 Webinar 1. Available: https:// www.youtube.com/watch?v=CjEhV68GcD8
- 28 Szerlip M, Anwaruddin S, Aronow HD, et al. Considerations for cardiac catheterization laboratory procedures during the COVID-19 pandemic perspectives from the Society for cardiovascular angiography and interventions emerging leader mentorship (Scal elm) members and graduates. *Catheter Cardiovasc Interv* 2020.
- 29 Tang N, Li D, Wang X, *et al*. Abnormal coagulation parameters are associated with poor prognosis in patients with novel coronavirus pneumonia. *J Thromb Haemost* 2020;18:844–7.
- 30 Sugiura M, Hiraoka K, Ohkawa S, et al. A clinicopathological study on cardiac lesions in 64 cases of disseminated intravascular coagulation. Jpn Heart J 1977;18:57–69.
- 31 Wang YD, Zhang SP, Wei QZ, et al. [COVID-19 complicated with DIC: 2 cases report and literatures review]. Zhonghua Xue Ye Xue Za Zhi 2020;41:E001.

Review

- 32 Fried JA, Ramasubbu K, Bhatt R, *et al*. The variety of cardiovascular presentations of COVID-19. *Circulation* 2020.
- 33 Sala S, Peretto G, Gramegna M, et al. Acute myocarditis presenting as a reverse Tako-Tsubo syndrome in a patient with SARS-CoV-2 respiratory infection. Eur Heart J 2020. doi:10.1093/eurhearti/ehaa286. [Epub ahead of print: 08 Apr 2020].
- 34 Arentz M, Yim E, Klaff L, et al. Characteristics and outcomes of 21 critically ill patients with COVID-19 in Washington state. JAMA 2020. doi:10.1001/jama.2020.4326. [Epub ahead of print: 19 Mar 2020].
- 35 Inciardi RM, Lupi L, Zaccone G, *et al*. Cardiac involvement in a patient with coronavirus disease 2019 (COVID-19). *JAMA Cardiol* 2020. doi:10.1001/ jamacardio.2020.1096. [Epub ahead of print: 27 Mar 2020].
- 36 Mehta P, McAuley DF, Brown M, et al. COVID-19: consider cytokine storm syndromes and immunosuppression. *Lancet* 2020;395:1033–4.
- 37 Wang Z, Zhang A, Wan Y, et al. Early hypercytokinemia is associated with interferoninduced transmembrane protein-3 dysfunction and predictive of fatal H7N9 infection. Proc Natl Acad Sci U S A 2014;111:769–74.
- 38 Wong CK, Lam CWK, Wu AKL, *et al.* Plasma inflammatory cytokines and chemokines in severe acute respiratory syndrome. *Clin Exp Immunol* 2004;136:95–103.
- 39 Kim ES, Choe PG, Park WB, et al. Clinical progression and cytokine profiles of middle East respiratory syndrome coronavirus infection. J Korean Med Sci 2016;31:1717–25.
- 40 Mann DL. Innate immunity and the failing heart: the cytokine hypothesis revisited. *Circ Res* 2015;116:1254–68.
- 41 Ruan Q, Yang K, Wang W, *et al*. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive Care Med* 2020.
- 42 Liu K, Fang Y-Y, Deng Y, et al. Clinical characteristics of novel coronavirus cases in tertiary hospitals in Hubei Province. *Chin Med J* 2020. doi:10.1097/ CM9.000000000000744. [Epub ahead of print: 07 Feb 2020].
- 43 Shi Z, Fu W. Diagnosis and treatment recommendation for novel coronavirus pneumonia related isolated distal deep vein thrombosis. *Shanghai Medical Journal*, 2020. Available: http://kns.cnki.net/kcms/detail/31.1366.R.20200225.1444.004.html
- 44 Klok FA, Kruip MJHA, van der Meer NJM, et al. Incidence of thrombotic complications in critically ill ICU patients with COVID-19. *Thromb Res* 2020. doi:10.1016/j. thromres.2020.04.013. [Epub ahead of print: 10 Apr 2020].
- 45 Han H, Yang L, Liu R, et al. Prominent changes in blood coagulation of patients with SARS-CoV-2 infection. *Clin Chem Lab Med* 2020. doi:10.1515/cclm-2020-0188. [Epub ahead of print: 16 Mar 2020].
- 46 Shanghai Clinical Treatment Expert Group for COVID-19. [Comprehensive treatment and management of coronavirus disease 2019: expert consensus statement from Shanghai]. Chin J Infect 2020.
- 47 Tang N, Bai H, Chen X, et al. Anticoagulant treatment is associated with decreased mortality in severe coronavirus disease 2019 patients with coagulopathy. J Thromb Haemost 2020. doi:10.1111/jth.14817. [Epub ahead of print: 27 Mar 2020].
- 48 Sommerstein RGC. Preventing a COVID-19 pandemic: ACE inhibitors as a potential risk factor for fatal COVID-19. BMJ 2020;368:M810-M.
- 49 Crackower MA, Sarao R, Oudit GY, *et al*. Angiotensin-converting enzyme 2 is an essential regulator of heart function. *Nature* 2002;417:822–8.
- 50 Yang P, Gu H, Zhao Z, *et al*. Angiotensin-converting enzyme 2 (ACE2) mediates influenza H7N9 virus-induced acute lung injury. *Sci Rep* 2014;4:7027.
- 51 Vaduganathan M, Vardeny O, Michel T, et al. Renin-Angiotensin-Aldosterone system inhibitors in patients with Covid-19. N Engl J Med 2020. doi:10.1056/ NEJMsr2005760. [Epub ahead of print: 30 Mar 2020].
- 52 Ferrario CM, Jessup J, Chappell MC, et al. Effect of angiotensin-converting enzyme inhibition and angiotensin II receptor blockers on cardiac angiotensin-converting enzyme 2. *Circulation* 2005;111:2605–10.
- 53 Luque M, Martin P, Martell N, *et al*. Effects of captopril related to increased levels of prostacyclin and angiotensin-(1-7) in essential hypertension. *J Hypertens* 1996;14:799–805.
- 54 Wang M, Cao R, Zhang L, *et al*. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res* 2020;30:269–71.
- 55 Sahraei Z, Shabani M, Shokouhi S, *et al*. Aminoquinolines against coronavirus disease 2019 (COVID-19): chloroquine or hydroxychloroquine. *Int J Antimicrob Agents* 2020;105945.
- 56 Gao J, Tian Z, Yang X. Breakthrough: chloroquine phosphate has shown apparent efficacy in treatment of COVID-19 associated pneumonia in clinical studies. *Biosci Trends* 2020;14:72–3.
- 57 Gautret P, Lagier J-C, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. Int J Antimicrob Agents 2020;105949:105949.
- Retallack H, Di Lullo E, Arias C, *et al.* Zika virus cell tropism in the developing human brain and inhibition by azithromycin. *Proc Natl Acad Sci U S A* 2016;113:14408–13.
 Simona C, Laguage C, Davier M, Davier M, Collog C, Simona C, Laguage C, Davier M, Davier M, Collog C, Simona C, Laguage C, Davier M, Collog C, Simona C, Laguage C, Davier M, Collog C, Simona C, Laguage C, Davier M, Collage C, Calabara C,
- Simoens S, Laekeman G, Decramer M. Preventing COPD exacerbations with macrolides: a review and budget impact analysis. *Respir Med* 2013;107:637–48.
 White NJ. Cardiotoxicity of antimalarial drugs. *Lancet Infect Dis* 2007;7:549–58.

- 61 Fohner AE, Sparreboom A, Altman RB, et al. PharmGKB summary: macrolide antibiotic pathway, pharmacokinetics/pharmacodynamics. *Pharmacogenet Genomics* 2017;27:164–7.
- 62 Kim K-A, Park J-Y, Lee J-S, et al. Cytochrome P450 2C8 and CYP3A4/5 are involved in chloroquine metabolism in human liver microsomes. Arch Pharm Res 2003;26:631–7.
- 63 Lakkireddy DR, Chung MK, Gopinathannair R, *et al*. Guidance for cardiac electrophysiology during the coronavirus (COVID-19) pandemic from the heart rhythm Society COVID-19 Task force; electrophysiology section of the American College of cardiology; and the electrocardiography and arrhythmias Committee of the Council on clinical cardiology, American heart association. *Circulation* 2020. doi:10.1161/CIRCULATIONAHA.120.047063. [Epub ahead of print: 31 Mar 2020].
- 64 Russell CD, Millar JE, Baillie JK. Clinical evidence does not support corticosteroid treatment for 2019-nCoV lung injury. *Lancet* 2020;395:473–5.
- 65 Fardet L, Galicier L, Lambotte O, *et al*. Development and validation of the HScore, a score for the diagnosis of reactive hemophagocytic syndrome. *Arthritis Rheumatol* 2014;66:2613–20.
- 66 Xu X, Han M, Li T, *et al.* Effective treatment of severe COVID-19 patients with tocilizumab. *ChinaXiv* 2020:20200300026.
- 67 National Cancer Institute Naples. Tocilizumab in COVID-19 pneumonia (TOCIVID-19). Available: https://www.clinicaltrials.gov/ct2/show/NCT04317092
- 68 Sanofi. Sarilumab COVID-19. Available: https://www.clinicaltrials.gov/ct2/show/ NCT04327388
- 69 Ramanathan K, Antognini D, Combes A, *et al*. Planning and provision of ECMO services for severe ARDS during the COVID-19 pandemic and other outbreaks of emerging infectious diseases. *Lancet Respir Med* 2020. doi:10.1016/S2213-2600(20)30121-1. [Epub ahead of print: 20 Mar 2020].
- 70 Ukimura A, Ooi Y, Kanzaki Y, *et al*. A national survey on myocarditis associated with influenza H1N1pdm2009 in the pandemic and postpandemic season in Japan. *J Infect Chemother* 2013;19:426–31.
- 71 Mertz D, Kim TH, Johnstone J, *et al*. Populations at risk for severe or complicated influenza illness: systematic review and meta-analysis. *BMJ* 2013;347:f5061.
- 72 Chan JWM, Ng CK, Chan YH, *et al.* Short term outcome and risk factors for adverse clinical outcomes in adults with severe acute respiratory syndrome (SARS). *Thorax* 2003;58:686–9.
- 73 Badawi A, Ryoo SG. Prevalence of comorbidities in the middle East respiratory syndrome coronavirus (MERS-CoV): a systematic review and meta-analysis. Int J Infect Dis 2016;49:129–33.
- 74 CDC. Preliminary estimates of the prevalence of selected underlying health conditions among patients with coronavirus disease 2019 — United States 2020.
- 75 Virani SS, Alonso A, Benjamin EJ, et al. Heart disease and stroke Statistics-2020 update: a report from the American heart association. *Circulation* 2020;141:e139–596.
- 76 Croft JB, Wheaton AG, Liu Y, et al. Urban-Rural County and State Differences in Chronic Obstructive Pulmonary Disease - United States, 2015. MMWR Morb Mortal Wkly Rep 2018;67:205–11.
- 77 Hu S, Gao R, Liu L, et al. Summary of the 2018 report on cardiovascular disease in China. Chinese Circulation Journal 2019;34.
- 78 Garg S, Kim L, Whitaker M, *et al.* Hospitalization Rates and Characteristics of Patients Hospitalized with Laboratory-Confirmed Coronavirus Disease 2019 - COVID-NET, 14 States, March 1-30, 2020. *MMWR Morb Mortal Wkly Rep* 2020;69:458–64.
- 79 Grasselli G, Zangrillo A, Zanella A, et al. Baseline characteristics and outcomes of 1591 patients infected with SARS-CoV-2 admitted to ICUs of the Lombardy region, Italy. JAMA 2020. doi:10.1001/jama.2020.5394. [Epub ahead of print: 06 Apr 2020].
- 80 CDC FluServ-Net. Weekly U.S. influenza surveillance report. Available: https://www.cdc.gov/flu/weekly/
- 81 Booth CM, Matukas LM, Tomlinson GA, et al. Clinical features and shortterm outcomes of 144 patients with SARS in the greater Toronto area. JAMA 2003;289:2801–9.
- 82 Xu Z, Shi L, Wang Y, et al. Pathological findings of COVID-19 associated with acute respiratory distress syndrome. *Lancet Respir Med* 2020;8:420–2.
- 83 Tian S, Xiong Y, Liu H, et al. Pathological study of the 2019 novel coronavirus disease (COVID-19) through post-mortem core biopsies. Preprints 2020.
- 84 Fox SE, Akmatbekov A, Harbert JL, et al. Pulmonary and cardiac pathology in Covid-19: the first autopsy series from new Orleans. medRxiv 2020:2020.04.06.20050575.
- 85 Lang Z-W, Zhang L-J, Zhang S-J, et al. A clinicopathological study of three cases of severe acute respiratory syndrome (SARS). Pathology 2003;35:526–31.
- 86 Chen C, Chen C, Yan JT, et al. [Analysis of myocardial injury in patients with COVID-19 and association between concomitant cardiovascular diseases and severity of COVID-19]. Zhonghua Xin Xue Guan Bing Za Zhi 2020;48:E008.
- 87 Zeng JHL, Yuan Y, Wang J, et al. First case of COVID-19 infection with fulminant myocarditis complication: case report and insights. Preprints, 2020.